



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

271. Proposed by E. B. ESCOTT, Ann Arbor, Mich.

In the differential equation

$$\left(\frac{d^2y}{dx^2}\right)^2 \frac{d^5y}{dx^5} - 5\left(\frac{d^2y}{dx^2}\right) \frac{d^3y}{dx^3} \frac{d^4y}{dx^4} + \frac{40}{9} \left(\frac{d^3y}{dx^3}\right)^2 = 0,$$

show that there is an integrating factor of the form $\left(\frac{d^2y}{dx^2}\right)^n$, and integrate the equation.

Solution by the PROPOSER, and LEVI S. SHIVELY, Mt. Morris College, Mt. Morris, Ill.

When the equation is multiplied by $(d^2y/dx^2)^n$, if it is an exact equation, its first integral is

$$\left(\frac{d^2y}{dx^2}\right)^{n+2} \frac{d^4y}{dx^4} + \frac{40}{9(n+1)} \left(\frac{d^2y}{dx^2}\right)^{n+1} \left(\frac{d^3y}{dx^3}\right)^2 = c_1 \dots (2).$$

Differentiating this equation and comparing with original equation, we find that $n = -\frac{1}{3}$ and $-\frac{1}{3}$.

In (2), put $n = -\frac{1}{3}$. The equation is exact and its integral is

$$\left(\frac{d^2y}{dx^2}\right)^{-\frac{2}{3}} \frac{d^3y}{dx^3} = c_1x + c_2.$$

Integrating again,

$$-\frac{3}{2} \left(\frac{d^2y}{dx^2}\right)^{-\frac{2}{3}} = \frac{c_1}{2}x^2 + c_2x + c_3.$$

Solving for d^2y/dx^2 , and changing the constants,

$$\frac{d^2y}{dx^2} = \frac{1}{(c_1x^2 + 2c_2x + c_3)^{\frac{3}{2}}}.$$

Integrating twice, we have

$$y = \frac{1}{c_1c_3 - c_2^2} \sqrt{(c_1x^2 + 2c_2x + c_3) + c_4x + c_5}.$$

Also solved by G. B. M. Zerr, and V. M. Spunar.

272. Proposed by CLARENCE OHLENDORF, Chicago, Ill.

Find $\int \log_e \tan^{-1} x dx$.

Solution by J. SCHEFFER, A. M., Hagerstown, Md., and C. N. SCHMALL, New York City.

Putting $x = \tan y$, we have

$$\int \log \tan^{-1} x dx = \int \log y \frac{dy}{\cos^2 y} = \int \log y \sec^2 y dy = \log y - \tan y - \int \frac{\tan y dy}{y}.$$

Substituting for $\tan y$ its series, and integrating, we get

$$\int \frac{\tan y dy}{y} = \frac{2^2(2^2-1)}{2!} B_1 y + \frac{2^4(2^4-1)}{4!} B_3 \frac{y^3}{3} + \frac{2^6(2^6-1)}{6!} B_5 \frac{y^5}{5},$$

where B_1, B_3, B_5, \dots are Bernoulli's numbers.

Therefore the given integral is

$$x \log \tan^{-1} x - \left[\frac{2^2(2^2-1)}{2!} B_1 \tan^{-1} x + \frac{2^4(2^4-1)}{4!} B_3 \frac{(\tan^{-1} x)^3}{3} + \dots \right].$$

Also solved by G. B. M. Zerr, V. M. Spunar, and J. E. Sanders.

273. Proposed by J. SCHEFFER, A. M., Hagerstown, Md.

On one side of a circular pond a feet in radius is a duck. On the diametrically opposite side of the pond is a dog. Both begin to swim at the same time, the duck swimming around the circumference of the pond at the rate of m feet a minute, the dog swimming directly towards the duck at the rate of n feet per minute. How far will the dog swim in overtaking the duck?

Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

If the line joining the positions of the dog and the duck is tangent to the dog's path, the following is the solution.

Let A, B be the starting points of dog and duck; O , the center of the pond; P, R , corresponding positions of the dog and duck. Then the tangent t , from P must pass through R . Let the duck swim to S and the dog to Q . Draw OF, PH perpendicular to AB , PI, QD perpendicular to OF , QC perpendicular to PI , and RE perpendicular to QS , produced.

Let $AB=2a$, $AP=\sigma$, $\angle RAB=\phi$, $OD=x$, $n/m=b$. Then $PR=d\sigma$, $\angle RAS=d\phi$, $DI=QC=dx$.

Now $\sigma=2ab\phi$; $\therefore d\sigma/2ad\phi=b$.

In the limit the triangles PQC and RES are similar.

$\therefore d\sigma : dx = 2ad\phi : ES$. $\therefore ES = (2ad\phi/d\sigma)dx = dx/b$.

The tangent t has negative increments at both ends (PQ and ES).

$\therefore dt = -d\sigma - dx/b$, or $t = C - \sigma - x/b$. When $t=2a$, $\sigma=0$, $x=0$.

$\therefore C=2a$. $\therefore t=2a-2ab\phi-x/b$. When $t=0$, $x=2a\sin\phi\cos\phi$.

$\therefore b=b^2\phi+\sin\phi\cos\phi=(b^2+1)\phi-\frac{2}{3}\phi^3+\frac{2}{15}\phi^5-\frac{4}{315}\phi^7+\dots$

$$Q = \frac{b}{b^2+1} + \frac{2b^4}{3(b^2+1)^4} + \frac{2b^5(9-b^2)}{15(b^2+1)^7} + \frac{4(225+b^4-54b^2)b^7}{315(b^2+1)^{10}} + \dots$$

by reversion of series.